

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Wood dust and urinary 15-F2t isoprostane in Italian industry workers

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1695536> since 2019-04-03T16:00:20Z

Published version:

DOI:10.1016/j.envres.2019.03.058

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Wood dust and urinary 15-F_{2t} isoprostane in Italian industry workers**

2 Roberto Bono^a, Fabio Capacci^b, Filippo Cellai^c, Carla Sgarrella^b, Valeria Bellisario^a, Giulia Trucco^a,
3 Lorenzo Tofani^d, Alessio Peluso^e, Carla Poli^f, Luciano Arena^f, Sara Piro^g, Lucia Miligi^g, Armelle
4 Munnia^c, and Marco Peluso^{c,*}

5 ^a *Department of Public Health and Pediatrics, University of Turin, Italy*

6 ^b *Functional Unit for Prevention, Health and Safety in the Workplace, ASL10, Florence, Italy*

7 ^c *Cancer Factor Risk Branch, Regional Cancer Prevention Laboratory, ISPRO-Study, Prevention
8 and Oncology Network Institute, 50139 - Florence, Italy*

9 ^d *Department of Neurosciences, Psychology, Drug Research and Child Health, University of
10 Florence, Florence, Italy*

11 ^e *Statistician, Florence, Italy*

12 ^f *Department of Prevention, ASL11, Empoli, Florence, Italy*

13 ^g *Unit of Environmental and Occupational Epidemiology, ISPRO-Study, Prevention and Oncology
14 Network Institute, 50139 - Florence, Italy*

15 *Correspondence to: Cancer Factor Risk Branch, Regional Cancer Prevention Laboratory, ISPRO-
16 Study, Prevention and Oncology Network Institute, 50139 - Florence, Italy. E-mail address:
17 m.peluso@ispro.toscana.it

18

Abstract

Wood dust is one of the most common occupational exposures, with about 3.6 million of workers in the wood industry in Europe. Wood particles can deposit in the nose and the respiratory tract and cause adverse health effects. Occupational exposure to wood dust has been associated with malignant tumors of the nasal cavity and paranasal sinuses. The induction of oxidative stress and the generation of reactive oxygen species through activation of inflammatory cells could have a role in the carcinogenicity of respirable wood dust. Therefore, we conducted a cross-sectional study to evaluate the prevalence of urinary 15-F_{2t} isoprostane (15-F_{2t}-IsoP), a biomarker of oxidative stress and peroxidation of lipids, in 123 wood workers compared to 57 unexposed controls living in Tuscany region, Italy. 15-F_{2t}-IsoP generation was measured by ELISA. The main result of the present study showed that a statistically significant excess of this biomarker occurred in the workers exposed to 1.48 mg/m³ of airborne wood dust with respect to the unexposed controls (0.05 mg/m³). The overall mean ratio (MR) between the workers exposed to wood dust and the controls was 1.36, 95% Confidence Interval (C.I.) 1.18–1.57, after correction for age and smoking habits. A significant increment of 15-F_{2t}-IsoP (43%) was observed in the smokers as compared to the non-smokers. The urinary excretion of 15-F_{2t}-IsoP was significantly associated with co-exposure to organic solvents and formaldehyde, i.e., MR of 1.41, 95% C.I. 1.17-1.70, after adjustment for age and smoking habits. A 41% excess was observed in long-term wood workers, 95% C.I. 1.14-1.75. Multivariate regression analysis showed that the level of 15-F_{2t}-IsoP was linearly correlated to the length of exposure, regression coefficient (β) = 0.244 \pm 0.002 (SE). The overall increment by exposure group persisted after stratification for smoking habits. For instance, in smokers, a 53% excess was detected in the wood workers as compared to the controls, 95% C.I. 1.23-1.91. Our data support the hypothesis that oxidative stress and lipid peroxidation can have a role in the toxicity of wood dust. F₂-IsoP measure can be a tool for the evaluation of the effectiveness of targeted interventions aimed to reduce exposures to environmental carcinogens.

Key words: wood dust, organic solvents, formaldehyde, 15-F_{2t} isoprostane, primary prevention, occupational health.

47 1. Introduction

48 Wood dust is one of the most common occupational exposures, with about 3.6 million of workers in
49 the wood industry in Europe (Kauppinen et al., 2006). Wood particles can deposit in the nose and
50 the respiratory tract and cause adverse health effects (Çelik and Kanık, 2006). Epidemiological
51 studies have indeed associated the exposure to wood dust to sinonasal cancers (SNC) (Acheson
52 et al., 1968; Ball, 1968). In 1960, the first association with SNC was shown in the wood industry
53 (Acheson et al., 1968). In 1995, this agent was classified as carcinogenic to humans (Group 1) by
54 the International Agency for Research on Cancer (IARC) based mostly on a SNC excess (IARC,
55 1995). In 2012, the IARC confirmed the human carcinogenicity of wood dust and reported the first
56 link with nasopharynx cancer (IARC, 2012). Considering other types of cancer, a meta-analysis
57 has suggested a relationship with lung cancer (Hancock et al., 2015), but a significant influence of
58 the geographic region was apparent.

59 SNC has been under compulsory surveillance since 2008 in Italy, through the “Sinonasal Cancer
60 National Registry” (Registro Nazionale Tumori Naso-Sinusali: ReNaTuNS), a nationwide cancer
61 registry coordinated by the National Institute for Insurance Against Accidents at Work (Istituto
62 Nazionale per l'Assicurazione contro gli Infortuni sul Lavoro: INAIL) (Binazzi et al., 2017).
63 Currently, the registry covers a proportion of Italy through regional structures devoted to the active
64 search for cases from hospitals, to the definition of the modalities of exposure and has recorded
65 1,529 cases between 2000-2016. A study conducted by Demers et al. (Demers et al., 1995) found
66 a doubled risk statistically significant for sinonasal cancer in men employed in any wood-related job
67 (OR = 2.0, 95% CI: 1.6 to 2.5) in comparison to men who had never worked in a wood-related job.
68 The increased risk was found among sawmill workers (OR = 2.5, 95% CI: 1.8 to 3.4), furniture
69 workers (OR = 4.5, 95% CI: 3.2 to 6.5) and carpenters (OR = 2.9, 95% CI: 2.1 to 3.9), while no
70 excess risk was observed among forestry, logging, pulp and paper workers. An increasing risk was
71 detected in relation to the duration of exposure, and lagging exposure by 5, 10 or 20 years
72 increased the strength of the association between duration of employment and sinonasal
73 adenocarcinoma. Elevated risk for adenocarcinoma of the nasal cavity and paranasal sinuses
74 (ADCN), a SNC subtype frequently associated with wood dust exposure (IARC, 2012), OR 58.6,
75 95% C.I. 23.74-144.8, was even reported among wood workers of the Piedmont region, Italy
76 (d'Errico et al., 2009). Stronger ADCN risk, OR 179.9, 95% C.I. 55.37-584.4, was found among
77 those workers exposed to high level of wood dust (d'Errico et al., 2009).

78 Higher levels of oxidative damage, measured by the micronucleus and the comet assays in blood,
79 buccal and nasal cells, have been detected in wood workers compared to unexposed controls
80 (Bruschweiler et al., 2016; Palus et al., 1999; Rekhadevi et al., 2009). An enhanced risk for
81 chromosomal instability was found in wood workers (Bruschweiler et al., 2014; Çelik and Kanık,
82 2006; Rekhadevi et al., 2009). Discrepant results have been reported (Wultsch et al., 2015). In that
83 study, no induction of micronuclei was observed in wood workers exposed to 0.39-0.66 mg/m³
84 wood dust levels. Thus, further investigation into wood workers' occupational exposures are
85 warranted. Furthermore, co-exposures to chrome, organic solvents, tannins, formaldehyde, textile
86 dust and pesticides have been reported in the wood industry (Binazzi et al., 2017). In 2012, IARC
87 suggested that the cancer risk of wood workers could be associated with the inflammatory
88 reactions following wood dust exposure rather than to the direct action of this carcinogen (IARC,
89 2012). Inflammatory cells can generate a large spectrum of proinflammatory mediators and free
90 radicals (Pylkkänen et al., 2009). Excessive production of reactive oxygen species (ROS) can
91 cause damage to lipids, proteins and DNA (Marnett, 2000). Peroxidation of lipids (LPO) can lead to
92 the production of aldehydes, such as malondialdehyde and 4-hydroxynonenal (Marnett, 2000), as
93 well as to secondary oxidation products such as a series of prostaglandin-like products termed
94 isoprostanes (IsoPs) (Roberts and Morrow, 2000).
95 IsoPs are compounds generated from the non-enzymatic free radical-catalyzed peroxidation of
96 arachidonic acid and other highly unsaturated polyunsaturated fatty acids (Janicka et al., 2010).
97 IsoPs can be grouped into 4 subfamilies, denoted as 5-, 12-, 8-, or 15-series regioisomers,
98 depending on the carbon atom to which the side chain hydroxyl is attached. Among the three major
99 classes of IsoPs (F₂-, D₂- and E₂-), F₂-IsoPs are recognized as the most suitable biomarker for

100 their chemical stability (Roberts and Morrow, 2000). The measurement of this biomarker is widely
101 used for the analysis of endogenous oxidative stress following ROS production and peroxidation of
102 lipids (Basu, 2008). F₂-IsoPs are more advantageous over other LPO biomarkers because they
103 can be detected in a variety of biological samples including plasma, urine, lavage fluid and red
104 blood cells (Milne et al., 2015). As IsoPs generate from LPO, their amounts provide an integrated
105 measurement of unbalanced oxidant-antioxidant status (Lowe et al., 2013; Montuschi et al., 2004).

106 In the current study, we have investigated the potential effects of occupational exposure to wood
107 dust in the wood product manufacturing sector in the Tuscany Region of Italy. A cross-sectional
108 study was conducted to analyze the concentration of a biomarker of oxidative stress and LPO (15-
109 F_{2t}-IsoP) in the workers exposed to wood dust. One of the main advantages of using biomarkers is
110 that one can study signals of carcinogen exposure without having to wait for health effects as in
111 classical epidemiological studies (Merlo et al., 1997; Munnia et al., 2017; Munnia et al., 2007;
112 Peluso et al., 1997; Peluso et al., 2012). Although F₂-IsoP can be evaluated in different biological
113 fluids, we employed urine due to its ready availability and the high stability of F₂-IsoP in this
114 medium (Morrow et al., 1999). Since obesity has been associated with increased F₂-IsoP
115 concentrations (Annor et al., 2017; Il'yasova et al., 2015), we have examined the relationships
116 between urinary F₂-IsoPs and weight gain. Further understanding of the link between wood dust
117 and oxidative stress will improve knowledge of the mechanisms of carcinogenicity of this
118 occupational agent. Novelty of the current study is based on various items, including larger sample
119 size, a different geographical area, and a different type of data, i.e., the measurement of F₂-IsoPs
120 in urine rather than of micronucleus and DNA strand-breaks in blood, buccal and nasal cells.

121 2. Material and methods

122 2.1 Subjects and sampling

123 A sample of 44 wood companies of the province of Florence, Tuscany, Italy was randomly selected
124 among those which are under compulsory health surveillance. Wood companies were contacted in
125 person by medical doctors with qualifications in occupational medicine. The inclusion criteria were
126 as follows: (a) only workers exposed to wood dust from wood industry; (b) only workers with a
127 minimal exposure time of 1 year; (c) only controls without occupational history in industries
128 entailing exposure to known or suspected carcinogens; and (d) only controls resident in areas with
129 no proximity to major air pollution sources. All the volunteers involved in the study live and work in
130 the province of Florence, Tuscany, Italy. A 15-F_{2t}-IsoP was determined using spot urine samples
131 collected in the morning at each workplace. Wood workers and the other subjects were contacted
132 by the local occupational health services. All the volunteers were informed about the study aim and
133 gave a written informed consent. A life-style questionnaire was filled by each participant (Peluso et
134 al., 2015). Detailed information on socio-demographic and anthropometric characteristics,
135 education level, exposure to active and passive tobacco smoke, occupational exposure to wood
136 dust, protective gear use, co-exposures to organic solvents, welding and motor exhaust fumes and
137 occupational history were obtained. Subjects who had never smoked were classified as non-
138 smokers, smokers who had quit smoking from at least one month prior were classified as ex-
139 smokers, while individuals who smoked at least one cigarette per day were classified as smokers.
140 The Body Mass Index (BMI) categories reported from the National Heart, Lung, and
141 Blood Institute (<https://www.nhlbi.nih.gov/>) were used for grouping the study participants in normal
142 weight persons (18.5-24.9 kg/m²), overweight persons (25-25.99 kg/m²) and obese persons (≥30
143 kg/m²). BMI was determined using self-reported weight and height. Study procedures were
144 performed in accordance with the Declaration of Helsinki for human studies and the guidelines of
145 the General Hospital Institutional Committee that reviewed and approved the present protocol.

146 2.2 Exposure data

147 Data on carcinogen exposure are collected by employers and regularly sent to the Italian Institute
148 for Occupational Safety and Prevention (ISPESL) (Italian legislative decree no. 626 of 19
149 September 1994). Such information is named exposure registries and includes quantitative
150 measurements of wood dust exposure. Companies are responsible for collecting the exposure

151 measurements in accordance with the EN 689:1995 regulation by the European Committee on
152 Standardization (Scarselli et al., 2008). For the purpose of this research, data on occupational
153 exposure measurements of wood dust recorded in the Information System for Recording
154 Occupational Exposures to Carcinogens (SIREP) were used to estimate environmental air
155 concentrations.

156 2.3 Urinary 15-F_{2t} isoprostane and creatinine measurement

157 The IsoP under investigation consists of one of the most abundant endogenous F₂-IsoPs, i.e., the
158 15-F_{2t}-IsoP, a biomarker considered to be representative for human oxidant status (Milne et al.,
159 2015), also referred to as 8-iso-prostaglandin F_{2α} (Roberts and Morrow, 2000). In the current
160 study, the concentrations of 15-F_{2t}-IsoP were analyzed using the competitive enzyme-linked
161 immunoassay (ELISA) with a specific microplate kit (Oxford, MI, USA), according to the
162 manufacturer's instructions, as previously reported (Bono et al., 2015; Romanazzi et al., 2013). In
163 order to normalize urinary dilution rate of 15-F_{2t}-IsoP an aliquot of urine was used to quantify the
164 concentration of creatinine by the kinetic Jaffé procedure (Bartels and Cikes, 1969).

165 2.4 Statistical analysis

166 The level of 15-F_{2t}-IsoP was expressed as ng/mg creatinine. Given the right-skewed distribution of
167 this biomarker, the data were log transformed to stabilize the variance and normalize the
168 distribution. Multivariate statistical analyses were applied using log-normal regression models
169 including age (continuous), tobacco smoking, i.e., non-smokers, ex-smokers, smokers,
170 occupational history (years), and BMI, as predictive variables to evaluate the association between
171 exposure to wood dust and the urinary excretion of 15-F_{2t}-IsoP in the study participants. **Results**
172 **were adjusted for age and smoking.** This was based on a previous study showing potential
173 associations between these variables and biomarker levels (Ceppi et al., 2011). Wood workers
174 were classified according to occupational exposures in two additional sub-groups: a) wood workers
175 exposed to wood dust alone and b) wood workers with co-exposures to organic solvents. The
176 regression parameters estimated from the models were interpreted as ratios [Means Ratio (MR)]
177 between the means of 15-F_{2t}-IsoPs of each level of the categorical variables with respect to the
178 reference level, as appropriate. The MR was used as a measure of effect (van Houwelingen et al.,
179 2002). A p-value of <0.05 (two-tailed) was considered significant. Data were analyzed using
180 SAS9.3 and SPSS 20.0 (IBM SPSS Statistics, New York, NY).

181 3. Results

182 3.1 Study population

183 The underlying basic population consisted of workers employed in the wood product manufacturing
184 sector of the province of Florence, Tuscany Region, Italy. 32 out of 44 consented to participate to
185 the study. Participation rates were ~95%. The concentration of 15-F_{2t}-IsoPs in the wood workers
186 was evaluated along with control subjects, i.e., 123 wood workers and 57 controls. All participants
187 were males with a mean age of 45.3 ± 0.85 years and 35% of which were smokers. In the current
188 study, the wood workers consisted of carpenters and joiners, wood processing-plant operators,
189 woodworking machine operators, wood products assemblers, manufacturing labourers, industrial
190 robot operators and other wood related workers. The use of the most common Personal Protective
191 Equipment (PPE) in woodworking, i.e., disposable respirators, was generally reported from
192 majority of the wood workers. Controls were living in residential areas with no proximity to major air
193 pollution sources. The two groups had similar demographic, anthropometric and life-style
194 characteristics. The mean age of the wood workers and the controls was not statistically different
195 (Table 1). The average values of BMI were similar among the two groups (Table 1). The frequency
196 of smokers was similar between the groups, i.e., 36% of the wood workers and 37% of the
197 controls, respectively. The distribution of subjects with respect to wood dust exposure with – out
198 co-exposures to other airborne carcinogens and smoking habits was reported in Table 2. Other
199 variables included length of employment and BMI groups (Tables 1-2).

200 3.2 Exposure data

201 The exposure measurement of wood dust air concentrations corresponds to a single value
202 assessed from several consecutive samples by fixed positions (Scarselli et al., 2008). Airborne
203 levels of industrial contaminants were quantified by daily mean concentration, i.e., 8-h time-
204 weighted average (TWA-8), of respirable wood dust among exposed workers. The mean level of
205 TWA-8 concentration of wood dust was 1.48 mg/m³ in wood workers.

206 3.3 Urinary 15-F_{2t} isoprostane level, smoking habits and occupational exposure

207 An increased amount of 15-F_{2t}-IsoP was found in the urine of wood workers as compared to the
208 controls (4.2 vs 2.9 ng/mg creatinine, Table 2). The multivariate analysis shows that the 36%
209 excess of 15-F_{2t}-IsoP of the wood workers was significantly higher as compared to the controls,
210 95% C.I. 1.18–1.57. Smokers had an average concentration of 15-F_{2t}-IsoP higher than ex-smokers
211 and non-smokers. A significant excess was found in the smokers in respect to the non-smokers,
212 95% Confidence Interval (C.I.) 1.23–1.66, after adjusting for age by statistical analysis.
213 Subsequently, the effect of co-exposures to other potential occupational carcinogens in the wood
214 industry on the level of 15-F_{2t}-IsoP was investigated. Therefore, workers were stratified into two
215 additional sub-groups: a) only wood dust exposed workers and b) mixed exposed workers. Table 2
216 indicates that the highest level of 15-F_{2t}-IsoP occurred in the wood workers who were co-exposed
217 to respirable organic solvents in respect to those who were only exposed to wood dust (4.5 and 4.0
218 ng/mg creatinine, respectively). After adjusting for age and smoking, the multivariate analysis
219 shows a 41% increment of 15-F_{2t}-IsoP, 95% C.I. 1.17–1.70, in the mixed exposed workers,
220 whereas a lower increment was observed in the only wood dust exposed workers, 95% C.I. 1.15–
221 1.56. When we considered occupational history, there was a greater production of 15-F_{2t}-IsoP in
222 the long-term wood workers (4.8 ng/mg creatinine of 15-F_{2t}-IsoP) compared to those with shorter
223 occupational history (3.2 ng/mg creatinine). A 41% excess of 15-F_{2t}-IsoP was observed in the
224 wood workers with longer occupational exposure times, 95% C.I. 1.14–1.75. Then, the excretion of
225 15-F_{2t}-IsoPs was found to be significantly correlated with the length of dust exposure (p-value =
226 0.007). Table 3 reports the mean concentrations of 15-F_{2t}-IsoP and MR and 95% C.I. by exposure
227 group and smoking stratification. The highest amount of 15-F_{2t}-IsoP was found in the wood
228 workers who were smokers, i.e., 5.0 ng/mg.

229 3.4 Urinary 15-F_{2t} isoprostane level and BMI groups

230 Since early studies have supported the hypothesis of a relationship between F₂-IsoP and weight
231 gain (Annor et al., 2017; Il'yasova et al., 2015), the association of this biomarker of oxidant status
232 with BMI was investigated. Study participants were divided by three BMI categories: a) normal
233 weight persons (18.5-24.9 kg/m²), b) overweight persons (25-25.99 kg/m²) and c) obese persons
234 (≥30 kg/m²) to evaluate the relationship of F₂-IsoP with increase in body weight that could result in
235 excessive fat accumulation. Table 2 shows that the mean concentrations of 15-F_{2t}-IsoP of obese
236 and overweight participants were higher than those with normal weight, but, no significant effect
237 was found.

238 4. Discussion

239 Wood processing causes small particles of wood dust to become suspended in the air. Workers
240 can inhale these particles, which can cause adverse health effects. The main result of this paper
241 showed that significantly enhanced level of F₂-IsoP occurred in the workers compared to the
242 unexposed controls. A 36% excess of 15-F_{2t}-IsoP levels was found in the wood workers as
243 compared with the unexposed controls. Furthermore, the significant excess of 15-F_{2t}-IsoP
244 persisted after smoking habit stratification. Among the wood workers, a 53% excess of 15-F_{2t}-IsoP
245 was found in the smokers, a 48% excess was observed in the ex-smokers and a 27% in the non-
246 smokers as compared to the appropriate controls. The urinary excretion of this biomarker was
247 significantly associated with other parameters, including smoking habits, co-exposure to other
248 airborne carcinogens and length of employment. In particular, multivariate regression analysis
249 showed that the level of 15-F_{2t}-IsoP was linearly correlated to the length of exposure. In

agreement with our findings, other studies have previously reported increased oxidative stress generation in relation to occupational exposure to wood dust (Bruschweiler et al., 2016; Palus et al., 1999; Rekhadevi et al., 2009). Our findings provide strengthening of the hypothesis that oxidative stress and LPO can have a main role in the toxicity of wood dust. The analysis of F₂-IsoP in urine could offer a unique noninvasive analytic tool to study the role of ROS in chronic occupational exposures. In the current case, the linkage between urinary 15-F_{2t}-IsoPs and wood dust can be due to an increased production of ROS caused by inflammation after exposure fine and abundant airborne dust created during wood manipulation, maintenance activities and cleaning equipment. Increased oxidative stress and LPO can be caused from the oxidative burst of activated macrophages and neutrophils, cells with a main role in phagocytosis and clearance of xenobiotic particles, and from increased inflammatory cytokines and activated leukocytes (Gungor et al., 2010; Vanhees et al., 2013). This is in keeping with the results of previous studies using a biomarker of oxidative DNA damage and LPO (Bonassi et al., 2017; Bono et al., 2016; Bono et al., 2010; Peluso et al., 2013; Peluso et al., 2010). In support of our hypothesis, free radicals produced through chronic inflammatory process and cancer disease have been implicated as the causal factor in the mutagenesis of the *tumor suppressor gene TP53* (Brancato et al., 2016; Perez-Escuredo et al., 2012).

Next, our study showed an empirical relationship between tobacco smoking and the urinary excretion of 15-F_{2t}-IsoP, possibly related to the inhalation exposure to carcinogens contained in tobacco smoke. A 43% increment of the level of 15-F_{2t}-IsoP was present in overall the smokers as compared to the non-smokers. This excess is commonly interpreted as an harmful oxidative stress (Basu, 2008). These findings were somewhat expected as active smokers inhale a broad range of airborne carcinogens (IARC, 2004). The involvement of altered oxidative stress-related mechanisms in tobacco smoke carcinogenesis is in line with previous studies using various biomarkers of oxidative stress and LPO (Munnia et al., 2004; Peluso et al., 2014; Romanazzi et al., 2013). Various groups have measured the concentrations of F₂-IsoP in biological fluids of smokers. The mean level of free and esterified F₂-IsoP in the urine and plasma of smokers have been found to be significantly elevated as compared to non-smokers (Lowe et al., 2013). For instance, a previous cross-sectional study conducted on workers employed in an industry of plastic laminates in Piedmont, Italy, founds that smoking habits were significantly associated with the urinary excretion of 15-F_{2t}-IsoP (Romanazzi et al., 2013). When the relationship of 15-F_{2t}-IsoP with BMI was investigated, we found that the levels of 15-F_{2t}-IsoP tended to increase with fat accumulations. The 42% of the obese subjects showed indeed higher excretion of 15-F_{2t}-IsoP in respect to those with normal weight. This is partially in keeping with a previous work of Annor et al. (Annor et al., 2017) on the risk of diabetes and weight gain. In that study, the 35% of the obese individuals showed greater levels of F₂-IsoPs as compared to the controls. Additional studies are necessary to understand if this biomarker can be used as measure of lifestyle habits and intervention targeted to obesity prevention.

The threshold exposure limit recommended by the Italian law is 5 mg/m³ (Legislative Decree No 66/2000). This value will remain until the 2020th, after the entry into force of the new threshold exposure limit of 3 mg/m³ for five years and thereafter of 2 mg/m³ (European Directive Decree No 2017/2398). In this context, the SIREP database aims to facilitate analysis of occupational exposure figures for carcinogenic agents. In our study, the average amount of wood dust concentrations experienced from the wood workers was lower than threshold exposure limit of 3 mg/m³ (i.e., 1.48 mg/m³). This result is consistent with that reported from a previous study of Scarselli et al. (Scarselli et al., 2008), where the mean concentrations of wood dust was of 1.44 mg/m³ for 1.181 companies in Italy. Although our static measurements of the concentrations of industrial contaminants by fixed positions provide evidence of wood workers' exposure via air, they are not well representative of individual exposures to wood dust due to spatial and temporal variations. Therefore, we could not assess the potential relationships of airborne measurements with biomarker urinary excretion in exposed workers.

The airborne wood-dust concentrations from exposure registries are commonly used for the purposes of hazard control, exposure surveillance and assessment of health risks (Kauppinen et

al., 2006). Nevertheless, a limitation of our study is that no data on the variability of wood dust concentrations within a facility were available. The bias due to the variability of airborne carcinogen levels in occupational settings is difficult to predict, but a large variation can be present in one spot of a factory versus another. There could be an underestimation of the exposure to wood dust associated to some woodworking operations. For instance, local exhaust ventilation is used widely with fixed woodworking machinery, but it is generally lacking for hand tools (Pisaniello et al., 1991). The effects of poor work practices, such as the use of compressed air for cleaning, the lack of local exhaust ventilation for hand tools, that are commonly associated to high exposure levels to wood dust (Alwis et al., 1999), could be missed. Variations in the use of PPE (Alwis et al., 1999) and in the effective application of WorkSafe procedures at work places could have influenced the personal levels of exposure to wood dust of our workers.

Our subsequent finding shows that the urinary excretion of 15-F_{2t}-IsoP in the workers exposed to wood dust can aggravate with co-exposure to other respiratory carcinogens. An excess of 41% was detected in the wood workers that were co-exposed to organic solvents compared to the controls. Conversely, a lower excess was determined in the only wood dust exposed workers. High biosynthesis of F₂-IsoP can be due to frequent free radical-catalyzed reactions induced by alterations of oxidative stress, antioxidant defence and inflammation especially caused by occupational exposures to complex mixtures of airborne carcinogens. This is consistent with a cross-sectional study of workers exposed to dust containing silica (Peluso et al., 2015). In this study, the amount of oxidative stress and LPO biomarker of the workers exposed to airborne silica dust was greater in the case of occupational co-exposures to organic solvents, welding and motor exhaust fumes. Constituents of organic solvents, such as benzene and formaldehyde can be involved in the generation of oxidative stress and ROS (Bono et al., 2016; Bono et al., 2010; Sorensen et al., 2003) and cause the production of 15-F_{2t}-IsoP determined in the workers exposed to wood dust. Our results suggest that the urinary level of F₂-IsoP resulting from exposures to airborne wood dust can be affected from concomitant carcinogen exposures. Levels of oxidative stress can increase with exposures to organic solvents (Salimi et al., 2017; Singh et al., 2010), leading to a greater imbalance between excessive ROS generation and their degradation by antioxidants. The induction of reactive species can increase damage to membrane lipids, cellular proteins and DNA.

A significant difference in the amount of 15-F_{2t}-IsoP was then observed among sub-groups of wood workers with different occupational history. The urinary excretion of this biomarker of oxidant status was significantly elevated in those subjects with longer exposure time. An 41% excess of 15-F_{2t}-IsoP was found in the long-term wood workers as compared to those with shorter exposures, used as the reference level. Multivariate regression analysis showed that the level of 15-F_{2t}-IsoP was significantly linearly correlated to the length of employment, in agreement with a previous study on asbestos workers (Yoshida et al., 2001). In that study, the generation of an urinary biomarker of oxidative stress correlated positively with the length of exposure. Rekhadevi et al. (Rekhadevi et al., 2009) have similarly found an association between length of occupational exposure and increase frequency of micronuclei. Taken together, the occurrence of elevated oxidative stress in long-term wood workers can be possibly due to chronic inflammatory conditions. Our study suggests that the measure of urinary F₂-IsoPs can serve as a biomarker for assessing occupational carcinogen exposure and improving workplace safety. Particular effort should be devoted to studying long term health effects of exposure to wood dust, such as SNC.

Particular effort should be devoted to study delayed reactions such as diseases that take a long time to develop, like SNC, that can be caused by long-term exposure to this carcinogenic agent.

5. Conclusions

Our study provides a valuable contribution to the issue of oxidative stress in woodworking. An excessive ROS generation was demonstrated in exposed workers. Furthermore, we showed that exposure to organic solvents can increase the levels of urinary biomarkers of oxidative stress in wood workers. Results provide a basis for worker surveillance in occupational settings. F₂-IsoP measure could be used for the evaluation of the effectiveness of targeted interventions aimed to

356 reduce exposures to various environmental carcinogens. A more effective control of occupational
357 health risks could decrease the incidence of illness at work and improve the health of the
358 workforce.

359 **Acknowledgments**

360 We are grateful to Dr. Dusca Bartoli, Dr. Giuseppe A. Farina, Dr. Tonina E. Iaia, Dr. Pierluigi Faina,
361 Dr. Luciano Monticelli for the assistances contacting with workers.

362 **Funding**

363 This work was supported by the Tuscany Region and the National Institute for Insurance Against
364 Accident at Work (INAIL).

365 **Declarations of interest**

366 None

367

368 **References**

- 369 Acheson, E. D., et al., 1968. Nasal cancer in woodworkers in the furniture industry. *Br Med J.* 2, 587-596.
- 370 Alwis, U., et al., 1999. Dust Exposures in the Wood Processing Industry. *Am Ind Hyg Ass J.* 60, 641-646.
- 371 Annor, F., et al., 2017. African Ancestry Gradient Is Associated with Lower Systemic F(2)-Isoprostane Levels. *Oxid Med Cell Longev.* 2017, 8319176-8319176.
- 372 Ball, M. J., 1968. Nasal cancer in woodworkers. *Br Med J.* 3, 253-253.
- 373 Bartels, H., Cikes, M., 1969. [Chromogens in the creatinine determination of Jaffe]. *Clin Chim Acta.* 26, 1-10.
- 374 Basu, S., 2008. F2-isoprostanes in human health and diseases: from molecular mechanisms to clinical implications. *Antiox Redox Sign.* 10, 1405-1434.
- 375 Binazzi, A., et al., 2017. Sinonasal cancer in the Italian national surveillance system: Epidemiology, occupation, and public health implications. *Am J Ind Med.* 1–12.
- 376 Bonassi, S., et al., 2017. 3-(2-deoxy- β -d-erythro-penta-furanosyl) pyrimido [1, 2- α] purin-10 (3H)-one deoxyguanosine adducts of workers exposed to asbestos fibers. *Toxic Lett.* 270, 1-7.
- 377 Bono, R., et al., 2016. Formaldehyde-induced toxicity in the nasal epithelia of workers of a plastic laminate plant. *Toxicol Res.* 5, 752-760.
- 378 Bono, R., et al., 2010. Malondialdehyde-deoxyguanosine adduct formation in workers of pathology wards: the role of air formaldehyde exposure. *Chem Res Toxicol.* 23, 1342-8.
- 379 Bono, R., et al., 2015. Urban air and tobacco smoke as conditions that increase the risk of oxidative stress and respiratory response in youth. *Environ Res.* 137, 141-146.
- 380 Brancato, B., et al., 2016. 8-Oxo-7, 8-dihydro-2-deoxyguanosine and other lesions along the coding strand of the exon 5 of the tumour suppressor gene P53 in a breast cancer case-control study. *DNA Res.* 23, 395-402.
- 381 Bruschweiler, D. E., et al., 2016. DNA Damage among Wood Workers Assessed with the Comet Assay. *Environ Health Insights.* 10, 105-112.
- 382 Bruschweiler, E. D., et al., 2014. Workers exposed to wood dust have an increased micronucleus frequency in nasal and buccal cells: results from a pilot study. *Mutagenesis.* 29, 201-207.
- 383 Çelik, A., Kanık, A., 2006. Genotoxicity of occupational exposure to wood dust: Micronucleus frequency and nuclear changes in exfoliated buccal mucosa cells. *Environ Mol Mutagen.* 47, 693-698.
- 384 Ceppi, M., et al., 2011. Study design and statistical analysis of data in human population studies with the micronucleus assay. *Mutagenesis.* 26, 247-252.
- 385 d'Errico, A., et al., 2009. A case-control study on occupational risk factors for sino-nasal cancer. *Occup Environ Med.* 66, 448-455.
- 386 Demers, P. A., et al., 1995. Wood dust and sino-nasal cancer: pooled reanalysis of twelve case-control studies. *Am J Ind Med.* 28, 151-66.
- 387 Gungor, N., et al., 2010. Transcriptional profiling of the acute pulmonary inflammatory response induced by LPS: role of neutrophils. *Respir Res.* 11, 24.
- 388 Hancock, D. G., et al., 2015. Wood dust exposure and lung cancer risk: a meta-analysis. *Occup Environ Med.* 72, 889-898.
- 389 IARC, 1995. Wood Dust and Formaldehyde. *IARC Monogr Eval Carcinog Risk Chem Hum.* 62, 35-215.
- 390 IARC, 2004. Tobacco smoke and involuntary smoking. *IARC Monogr Eval Carcinog Risks Hum.* 83, 1-1438.
- 391 IARC, 2012. Wood Dust. A Review of Human Carcinogens: Arsenic, Metals, Fibres, and Dusts. *IARC Monogr Eval Carcinog Risks Hum.* 100 C, 1-469.
- 392 Il'yasova, D., et al., 2015. Urinary F2-Isoprostanes and Metabolic Markers of Fat Oxidation. *Oxid Medic Cell Long.* 2015.
- 393 Janicka, M., et al., 2010. Isoprostanes-Biomarkers of Lipid Peroxidation: Their Utility in Evaluating Oxidative Stress and Analysis. *IJMS.* 11, 4631.
- 394 Kauppinen, T., et al., 2006. Occupational Exposure to Inhalable Wood Dust in the Member States of the European Union. *Ann Occ Hyg.* 50, 549-561.
- 395 Lowe, F. J., et al., 2013. Lung cancer biomarkers for the assessment of modified risk tobacco products: an oxidative stress perspective. *Biomarkers.* 18, 183-195.
- 396 Marnett, L. J., 2000. Oxyradicals and DNA damage. *Carcinogenesis.* 21, 361-70.

419 Merlo, F., et al., 1997. Airborne levels of polycyclic aromatic hydrocarbons: 32P-postlabeling DNA adducts
420 and micronuclei in white blood cells from traffic police workers and urban residents. *J Environ*
421 *Pathol Toxicol Oncol.* 16, 157-62.

422 Milne, L. G., et al., 2015. The isoprostanes—25 years later. *BBA-Mol Cell Biol L.* 1851, 433-445.

423 Montuschi, P., et al., 2004. Isoprostanes: markers and mediators of oxidative stress. *FASEB.* 18, 1791-1800.

424 Morrow, J. D., et al., 1999. Quantification of the Major Urinary Metabolite of 15-F₂t-Isoprostane (8-iso-
425 PGF₂±) by a Stable Isotope Dilution Mass Spectrometric Assay. *Anal Bioch.* 269, 326-331.

426 Munnia, A., et al., 2004. Exocyclic malondialdehyde and aromatic DNA adducts in larynx tissues. *Free Radic*
427 *Biol Med.* 37, 850-8.

428 Munnia, A., et al., 2017. Bulky DNA Adducts, Tobacco Smoking, Genetic Susceptibility, and Lung Cancer
429 Risk. *Adv Clin Chem.* 81, 231-77.

430 Munnia, A., et al., 2007. 32P-Post-labelling method improvements for aromatic compound-related
431 molecular epidemiology studies. *Mutagenesis.* 22, 381-5.

432 Palus, J., et al., 1999. DNA damage detected by the comet assay in the white blood cells of workers in a
433 wooden furniture plant. *Mutat Res.* 444, 61-74.

434 Peluso, M., et al., 1997. Detection of DNA adducts in human nasal mucosa tissue by 32P-postlabeling
435 analysis. *Carcinogenesis.* 18, 339-44.

436 Peluso, M., et al., 2012. DNA methylation differences in exposed workers and nearby residents of the Ma Ta
437 Phut industrial estate, Rayong, Thailand. *Int J Epidemiol.* 41, 1753-60; author response 1761-3.

438 Peluso, M., et al., 2013. Malondialdehyde-deoxyguanosine and bulky DNA adducts in schoolchildren
439 resident in the proximity of the Sarroch industrial estate on Sardinia Island, Italy. *Mutagenesis.* 28,
440 315-21.

441 Peluso, M., et al., 2010. Malondialdehyde-deoxyguanosine adducts among workers of a Thai industrial
442 estate and nearby residents. *Environ Health Perspect.* 118, 55-9.

443 Peluso, M. E., et al., 2014. Aberrant methylation of hypermethylated-in-cancer-1 and exocyclic DNA
444 adducts in tobacco smokers. *Toxicol Sci.* 137, 47-54.

445 Peluso, M. E., et al., 2015. Oxidatively damaged DNA in the nasal epithelium of workers occupationally
446 exposed to silica dust in Tuscany region, Italy. *Mutagenesis.* 30, 519-25.

447 Perez-Escuredo, J., et al., 2012. Wood dust-related mutational profile of TP53 in intestinal-type sinonasal
448 adenocarcinoma. *Human Pathology.* 43, 1894-1901.

449 Pisaniello, D. L., et al., 1991. Wood dust exposure during furniture manufacture -results from an Australian
450 survey and consideration for threshold limit value development. *Am Ind Hyg Ass J.* 52, 485-492.

451 Pylkkänen, L., et al., 2009. Wood dusts induce the production of reactive oxygen species and caspase-3
452 activity in human bronchial epithelial cells. *Toxic in Vitro.* 262, 265-270.

453 Rekhadevi, P. V., et al., 2009. Genetic damage in wood dust-exposed workers. *Mutagenesis.* 24, 59-65.

454 Roberts, L. J., Morrow, J. D., 2000. Measurement of F₂-isoprostanes as an index of oxidative stress in vivo.
455 *Free Radic Biol Med.* 28, 505-513.

456 Romanazzi, V., et al., 2013. 15-F₂t isoprostane as biomarker of oxidative stress induced by tobacco smoke
457 and occupational exposure to formaldehyde in workers of plastic laminates. *Sci Total Environ.* 442,
458 20-5.

459 Salimi, A., et al., 2017. Xylene Induces Oxidative Stress and Mitochondria Damage in Isolated Human
460 Lymphocytes. *Toxic res.* 33, 233-238.

461 Scarselli, A., et al., 2008. Occupational exposure levels to wood dust in Italy, 1996–2006. *Occup*
462 *Environ Med.* 65, 567-574.

463 Singh, M. P., et al., 2010. Effects of co-exposure of benzene, toluene and xylene to *Drosophila*
464 *melanogaster*: Alteration in hsp70, hsp60, hsp83, hsp26, ROS generation and oxidative stress
465 markers. *Chemosphere.* 79, 577-587.

466 Sorensen, M., et al., 2003. Linking exposure to environmental pollutants with biological effects. *Mutat Res.*
467 544, 255-71.

468 van Houwelingen, H. C., et al., 2002. Advanced methods in meta-analysis: multivariate approach and meta-
469 regression. *Stat Med.* 21, 589-624.

470 Vanhees, K., et al., 2013. Intrauterine exposure to flavonoids modifies antioxidant status at adulthood and
471 decreases oxidative stress-induced DNA damage. *Free Radic Biol Med.* 57, 154-61.
472 Wultsch, G., et al., 2015. Impact of exposure to wood dust on genotoxicity and cytotoxicity in exfoliated
473 buccal and nasal cells. *Mutagenesis.* 30, 701-709.
474 Yoshida, R., et al., 2001. Urinary 8-oxo-7, 8-dihydro-2'-deoxyguanosine and biopyrrins levels among
475 construction workers with asbestos exposure history. *Ind Health.* 39, 186-8.
476
477

Wood dust and urinary 15-F_{2t} isoprostane in Italian industry workers

Roberto Bono^a, Fabio Capacci^b, Filippo Cellai^c, Carla Sgarrella^b, Valeria Bellisario^a, Giulia Trucco^a, Lorenzo Tofani^d, Alessio Peluso^e, Carla Poli^f, Luciano Arena^f, Sara Piro^g, Lucia Miligi^g, Armelle Munnia^c, and Marco Peluso^{c,*}

^a Department of Public Health and Pediatrics, University of Turin, Italy

^b Functional Unit for Prevention, Health and Safety in the Workplace, ASL10, Florence, Italy

^c Cancer Factor Risk Branch, Regional Cancer Prevention Laboratory, ISPRO-Study, Prevention and Oncology Network Institute, 50139 - Florence, Italy

^d Department of Neurosciences, Psychology, Drug Research and Child Health, University of Florence, Florence, Italy

^e Statistician, Florence, Italy

^f Department of Prevention, ASL11, Empoli, Florence, Italy

^g Unit of Environmental and Occupational Epidemiology, ISPRO-Study, Prevention and Oncology Network Institute, 50139 - Florence, Italy

*Correspondence to: Cancer Factor Risk Branch, Regional Cancer Prevention Laboratory, ISPRO-Study, Prevention and Oncology Network Institute, 50139 - Florence, Italy. E-mail address: m.peluso@ispro.toscana.it

Abstract

Wood dust is one of the most common occupational exposures, with about 3.6 million of workers in the wood industry in Europe. Wood particles can deposit in the nose and the respiratory tract and cause adverse health effects. Occupational exposure to wood dust has been associated with malignant tumors of the nasal cavity and paranasal sinuses. The induction of oxidative stress and the generation of reactive oxygen species through activation of inflammatory cells could have a role in the carcinogenicity of respirable wood dust. Therefore, we conducted a cross-sectional study to evaluate the prevalence of urinary 15-F_{2t} isoprostane (15-F_{2t}-IsoP), a biomarker of oxidative stress and peroxidation of lipids, in 123 wood workers compared to 57 unexposed controls living in Tuscany region, Italy. 15-F_{2t}-IsoP generation was measured by ELISA. The main result of the present study showed that a statistically significant excess of this biomarker occurred in the workers exposed to 1.48 mg/m³ of airborne wood dust with respect to the unexposed controls (0.05 mg/m³). The overall mean ratio (MR) between the workers exposed to wood dust and the controls was 1.36, 95% Confidence Interval (C.I.) 1.18–1.57, after correction for age and smoking habits. A significant increment of 15-F_{2t}-IsoP (43%) was observed in the smokers as compared to the non-smokers. The urinary excretion of 15-F_{2t}-IsoP was significantly associated with co-exposure to organic solvents and formaldehyde, i.e., MR of 1.41, 95% C.I. 1.17-1.70, after adjustment for age and smoking habits. A 41% excess was observed in long-term wood workers, 95% C.I. 1.14-1.75. Multivariate regression analysis showed that the level of 15-F_{2t}-IsoP was linearly correlated to the length of exposure, regression coefficient (β) = 0.244 \pm 0.002 (SE). The overall increment by exposure group persisted after stratification for smoking habits. For instance, in smokers, a 53% excess was detected in the wood workers as compared to the controls, 95% C.I. 1.23-1.91. Our data support the hypothesis that oxidative stress and lipid peroxidation can have a role in the toxicity of wood dust. F₂-IsoP measure can be a tool for the evaluation of the effectiveness of targeted interventions aimed to reduce exposures to environmental carcinogens.

Key words: wood dust, organic solvents, formaldehyde, 15-F_{2t} isoprostane, primary prevention, occupational health.

1. Introduction

Wood dust is one of the most common occupational exposures, with about 3.6 million of workers in the wood industry in Europe (Kauppinen et al., 2006). Wood particles can deposit in the nose and the respiratory tract and cause adverse health effects (Çelik and Kanık, 2006). Epidemiological studies have indeed associated the exposure to wood dust to sinonasal cancers (SNC) (Acheson et al., 1968; Ball, 1968). In 1960, the first association with SNC was shown in the wood industry (Acheson et al., 1968). In 1995, this agent was classified as carcinogenic to humans (Group 1) by the International Agency for Research on Cancer (IARC) based mostly on a SNC excess (IARC, 1995). In 2012, the IARC confirmed the human carcinogenicity of wood dust and reported the first link with nasopharynx cancer (IARC, 2012). Considering other types of cancer, a meta-analysis has suggested a relationship with lung cancer (Hancock et al., 2015), but a significant influence of the geographic region was apparent.

SNC has been under compulsory surveillance since 2008 in Italy, through the “Sinonasal Cancer National Registry” (Registro Nazionale Tumori Naso-Sinusali: ReNaTuNS), a nationwide cancer registry coordinated by the National Institute for Insurance Against Accidents at Work (Istituto Nazionale per l'Assicurazione contro gli Infortuni sul Lavoro: INAIL) (Binazzi et al., 2017). Currently, the registry covers a proportion of Italy through regional structures devoted to the active search for cases from hospitals, to the definition of the modalities of exposure and has recorded 1,529 cases between 2000-2016. A study conducted by Demers et al. (Demers et al., 1995) found a doubled risk statistically significant for sinonasal cancer in men employed in any wood-related job (OR = 2.0, 95% CI: 1.6 to 2.5) in comparison to men who had never worked in a wood-related job. The increased risk was found among sawmill workers (OR = 2.5, 95% CI: 1.8 to 3.4), furniture workers (OR = 4.5, 95% CI: 3.2 to 6.5) and carpenters (OR = 2.9, 95% CI: 2.1 to 3.9), while no excess risk was observed among forestry, logging, pulp and paper workers. An increasing risk was detected in relation to the duration of exposure, and lagging exposure by 5, 10 or 20 years increased the strength of the association between duration of employment and sinonasal adenocarcinoma. Elevated risk for adenocarcinoma of the nasal cavity and paranasal sinuses (ADCN), a SNC subtype frequently associated with wood dust exposure (IARC, 2012), OR 58.6, 95% C.I. 23.74-144.8, was even reported among wood workers of the Piedmont region, Italy (d'Errico et al., 2009). Stronger ADCN risk, OR 179.9, 95% C.I. 55.37-584.4, was found among those workers exposed to high level of wood dust (d'Errico et al., 2009).

Higher levels of oxidative damage, measured by the micronucleus and the comet assays in blood, buccal and nasal cells, have been detected in wood workers compared to unexposed controls (Bruschweiler et al., 2016; Palus et al., 1999; Rekhadevi et al., 2009). An enhanced risk for chromosomal instability was found in wood workers (Bruschweiler et al., 2014; Çelik and Kanık, 2006; Rekhadevi et al., 2009). Discrepant results have been reported (Wultsch et al., 2015). In that study, no induction of micronuclei was observed in wood workers exposed to 0.39-0.66 mg/m³ wood dust levels. Thus, further investigation into wood workers' occupational exposures are warranted. Furthermore, co-exposures to chrome, organic solvents, tannins, formaldehyde, textile dust and pesticides have been reported in the wood industry (Binazzi et al., 2017). In 2012, IARC suggested that the cancer risk of wood workers could be associated with the inflammatory reactions following wood dust exposure rather than to the direct action of this carcinogen (IARC, 2012). Inflammatory cells can generate a large spectrum of proinflammatory mediators and free radicals (Pylkkänen et al., 2009). Excessive production of reactive oxygen species (ROS) can cause damage to lipids, proteins and DNA (Marnett, 2000). Peroxidation of lipids (LPO) can lead to the production of aldehydes, such as malondialdehyde and 4-hydroxynonenal (Marnett, 2000), as well as to secondary oxidation products such as a series of prostaglandin-like products termed isoprostanes (IsoPs) (Roberts and Morrow, 2000).

IsoPs are compounds generated from the non-enzymatic free radical-catalyzed peroxidation of arachidonic acid and other highly unsaturated polyunsaturated fatty acids (Janicka et al., 2010). IsoPs can be grouped into 4 subfamilies, denoted as 5-, 12-, 8-, or 15-series regioisomers, depending on the carbon atom to which the side chain hydroxyl is attached. Among the three major classes of IsoPs (F₂-, D₂- and E₂-), F₂-IsoPs are recognized as the most suitable biomarker for

their chemical stability (Roberts and Morrow, 2000). The measurement of this biomarker is widely used for the analysis of endogenous oxidative stress following ROS production and peroxidation of lipids (Basu, 2008). F₂-IsoPs are more advantageous over other LPO biomarkers because they can be detected in a variety of biological samples including plasma, urine, lavage fluid and red blood cells (Milne et al., 2015). As IsoPs generate from LPO, their amounts provide an integrated measurement of unbalanced oxidant-antioxidant status (Lowe et al., 2013; Montuschi et al., 2004).

In the current study, we have investigated the potential effects of occupational exposure to wood dust in the wood product manufacturing sector in the Tuscany Region of Italy. A cross-sectional study was conducted to analyze the concentration of a biomarker of oxidative stress and LPO (15-F_{2t}-IsoP) in the workers exposed to wood dust. One of the main advantages of using biomarkers is that one can study signals of carcinogen exposure without having to wait for health effects as in classical epidemiological studies (Merlo et al., 1997; Munnia et al., 2017; Munnia et al., 2007; Peluso et al., 1997; Peluso et al., 2012). Although F₂-IsoP can be evaluated in different biological fluids, we employed urine due to its ready availability and the high stability of F₂-IsoP in this medium (Morrow et al., 1999). Since obesity has been associated with increased F₂-IsoP concentrations (Annor et al., 2017; Il'yasova et al., 2015), we have examined the relationships between urinary F₂-IsoPs and weight gain. Further understanding of the link between wood dust and oxidative stress will improve knowledge of the mechanisms of carcinogenicity of this occupational agent. Novelty of the current study is based on various items, including larger sample size, a different geographical area, and a different type of data, i.e., the measurement of F₂-IsoPs in urine rather than of micronucleus and DNA strand-breaks in blood, buccal and nasal cells.

2. Material and methods

2.1 Subjects and sampling

A sample of 44 wood companies of the province of Florence, Tuscany, Italy was randomly selected among those which are under compulsory health surveillance. Wood companies were contacted in person by medical doctors with qualifications in occupational medicine. The inclusion criteria were as follows: (a) only workers exposed to wood dust from wood industry; (b) only workers with a minimal exposure time of 1 year; (c) only controls without occupational history in industries entailing exposure to known or suspected carcinogens; and (d) only controls resident in areas with no proximity to major air pollution sources. All the volunteers involved in the study live and work in the province of Florence, Tuscany, Italy. A 15-F_{2t}-IsoP was determined using spot urine samples collected in the morning at each workplace. Wood workers and the other subjects were contacted by the local occupational health services. All the volunteers were informed about the study aim and gave a written informed consent. A life-style questionnaire was filled by each participant (Peluso et al., 2015). Detailed information on socio-demographic and anthropometric characteristics, education level, exposure to active and passive tobacco smoke, occupational exposure to wood dust, protective gear use, co-exposures to organic solvents, welding and motor exhaust fumes and occupational history were obtained. Subjects who had never smoked were classified as non-smokers, smokers who had quit smoking from at least one month prior were classified as ex-smokers, while individuals who smoked at least one cigarette per day were classified as smokers. The Body Mass Index (BMI) categories reported from the National Heart Lung, and Blood Institute (<https://www.nhlbi.nih.gov/>) were used for grouping the study participants in normal weight persons (18.5-24.9 kg/m²), overweight persons (25-25.99 kg/m²) and obese persons (≥30 kg/m²). BMI was determined using self-reported weight and height. Study procedures were performed in accordance with the Declaration of Helsinki for human studies and the guidelines of the General Hospital Institutional Committee that reviewed and approved the present protocol.

2.2 Exposure data

Data on carcinogen exposure are collected by employers and regularly sent to the Italian Institute for Occupational Safety and Prevention (ISPESL) (Italian legislative decree no. 626 of 19 September 1994). Such information is named exposure registries and includes quantitative measurements of wood dust exposure. Companies are responsible for collecting the exposure

151 measurements in accordance with the EN 689:1995 regulation by the European Committee on
152 Standardization (Scarselli et al., 2008). For the purpose of this research, data on occupational
153 exposure measurements of wood dust recorded in the Information System for Recording
154 Occupational Exposures to Carcinogens (SIREP) were used to estimate environmental air
155 concentrations.

156 2.3 Urinary 15-F_{2t} isoprostane and creatinine measurement

157 The IsoP under investigation consists of one of the most abundant endogenous F₂-IsoPs, i.e., the
158 15-F_{2t}-IsoP, a biomarker considered to be representative for human oxidant status (Milne et al.,
159 2015), also referred to as 8-iso-prostaglandin F_{2α} (Roberts and Morrow, 2000). In the current
160 study, the concentrations of 15-F_{2t}-IsoP were analyzed using the competitive enzyme-linked
161 immunoassay (ELISA) with a specific microplate kit (Oxford, MI, USA), according to the
162 manufacturer's instructions, as previously reported (Bono et al., 2015; Romanazzi et al., 2013). In
163 order to normalize urinary dilution rate of 15-F_{2t}-IsoP an aliquot of urine was used to quantify the
164 concentration of creatinine by the kinetic Jaffé procedure (Bartels and Cikes, 1969).

165 2.4 Statistical analysis

166 The level of 15-F_{2t}-IsoP was expressed as ng/mg creatinine. Given the right-skewed distribution of
167 this biomarker, the data were log transformed to stabilize the variance and normalize the
168 distribution. Multivariate statistical analyses were applied using log-normal regression models
169 including age (continuous), tobacco smoking, i.e., non-smokers, ex-smokers, smokers,
170 occupational history (years), and BMI, as predictive variables to evaluate the association between
171 exposure to wood dust and the urinary excretion of 15-F_{2t}-IsoP in the study participants. Results
172 were adjusted for age and smoking. This was based on a previous study showing potential
173 associations between these variables and biomarker levels (Ceppi et al., 2011). Wood workers
174 were classified according to occupational exposures in two additional sub-groups: a) wood workers
175 exposed to wood dust alone and b) wood workers with co-exposures to organic solvents. The
176 regression parameters estimated from the models were interpreted as ratios [Means Ratio (MR)]
177 between the means of 15-F_{2t}-IsoPs of each level of the categorical variables with respect to the
178 reference level, as appropriate. The MR was used as a measure of effect (van Houwelingen et al.,
179 2002). A p-value of <0.05 (two-tailed) was considered significant. Data were analyzed using
180 SAS9.3 and SPSS 20.0 (IBM SPSS Statistics, New York, NY).

181 3. Results

182 3.1 Study population

183 The underlying basic population consisted of workers employed in the wood product manufacturing
184 sector of the province of Florence, Tuscany Region, Italy. 32 out of 44 consented to participate to
185 the study. Participation rates were ~95%. The concentration of 15-F_{2t}-IsoPs in the wood workers
186 was evaluated along with control subjects, i.e., 123 wood workers and 57 controls. All participants
187 were males with a mean age of 45.3 ± 0.85 years and 35% of which were smokers. In the current
188 study, the wood workers consisted of carpenters and joiners, wood processing-plant operators,
189 woodworking machine operators, wood products assemblers, manufacturing labourers, industrial
190 robot operators and other wood related workers. The use of the most common Personal Protective
191 Equipment (PPE) in woodworking, i.e., disposable respirators, was generally reported from
192 majority of the wood workers. Controls were living in residential areas with no proximity to major air
193 pollution sources. The two groups had similar demographic, anthropometric and life-style
194 characteristics. The mean age of the wood workers and the controls was not statistically different
195 (Table 1). The average values of BMI were similar among the two groups (Table 1). The frequency
196 of smokers was similar between the groups, i.e., 36% of the wood workers and 37% of the
197 controls, respectively. The distribution of subjects with respect to wood dust exposure with – out
198 co-exposures to other airborne carcinogens and smoking habits was reported in Table 2. Other
199 variables included length of employment and BMI groups (Tables 1-2).

3.2 Exposure data

The exposure measurement of wood dust air concentrations corresponds to a single value assessed from several consecutive samples by fixed positions (Scarselli et al., 2008). Airborne levels of industrial contaminants were quantified by daily mean concentration, i.e., 8-h time-weighted average (TWA-8), of respirable wood dust among exposed workers. The mean level of TWA-8 concentration of wood dust was 1.48 mg/m³ in wood workers.

3.3 Urinary 15-F_{2t} isoprostane level, smoking habits and occupational exposure

An increased amount of 15-F_{2t}-IsoP was found in the urine of wood workers as compared to the controls (4.2 vs 2.9 ng/mg creatinine, Table 2). The multivariate analysis shows that the 36% excess of 15-F_{2t}-IsoP of the wood workers was significantly higher as compared to the controls, 95% C.I. 1.18–1.57. Smokers had an average concentration of 15-F_{2t}-IsoP higher than ex-smokers and non-smokers. A significant excess was found in the smokers in respect to the non-smokers, 95% Confidence Interval (C.I.) 1.23–1.66, after adjusting for age by statistical analysis. Subsequently, the effect of co-exposures to other potential occupational carcinogens in the wood industry on the level of 15-F_{2t}-IsoP was investigated. Therefore, workers were stratified into two additional sub-groups: a) only wood dust exposed workers and b) mixed exposed workers. Table 2 indicates that the highest level of 15-F_{2t}-IsoP occurred in the wood workers who were co-exposed to respirable organic solvents in respect to those who were only exposed to wood dust (4.5 and 4.0 ng/mg creatinine, respectively). After adjusting for age and smoking, the multivariate analysis shows a 41% increment of 15-F_{2t}-IsoP, 95% C.I. 1.17–1.70, in the mixed exposed workers, whereas a lower increment was observed in the only wood dust exposed workers, 95% C.I. 1.15–1.56. When we considered occupational history, there was a greater production of 15-F_{2t}-IsoP in the long-term wood workers (4.8 ng/mg creatinine of 15-F_{2t}-IsoP) compared to those with shorter occupational history (3.2 ng/mg creatinine). A 41% excess of 15-F_{2t}-IsoP was observed in the wood workers with longer occupational exposure times, 95% C.I. 1.14–1.75. Then, the excretion of 15-F_{2t}-IsoPs was found to be significantly correlated with the length of dust exposure (p-value = 0.007). Table 3 reports the mean concentrations of 15-F_{2t}-IsoP and MR and 95% C.I. by exposure group and smoking stratification. The highest amount of 15-F_{2t}-IsoP was found in the wood workers who were smokers, i.e., 5.0 ng/mg.

3.4 Urinary 15-F_{2t} isoprostane level and BMI groups

Since early studies have supported the hypothesis of a relationship between F₂-IsoP and weight gain (Annor et al., 2017; Il'yasova et al., 2015), the association of this biomarker of oxidant status with BMI was investigated. Study participants were divided by three BMI categories: a) normal weight persons (18.5-24.9 kg/m²), b) overweight persons (25-25.99 kg/m²) and c) obese persons (≥30 kg/m²) to evaluate the relationship of F₂-IsoP with increase in body weight that could result in excessive fat accumulation. Table 2 shows that the mean concentrations of 15-F_{2t}-IsoP of obese and overweight participants were higher than those with normal weight, but, no significant effect was found.

4. Discussion

Wood processing causes small particles of wood dust to become suspended in the air. Workers can inhale these particles, which can cause adverse health effects. The main result of this paper showed that significantly enhanced level of F₂-IsoP occurred in the workers compared to the unexposed controls. A 36% excess of 15-F_{2t}-IsoP levels was found in the wood workers as compared with the unexposed controls. Furthermore, the significant excess of 15-F_{2t}-IsoP persisted after smoking habit stratification. Among the wood workers, a 53% excess of 15-F_{2t}-IsoP was found in the smokers, a 48% excess was observed in the ex-smokers and a 27% in the non-smokers as compared to the appropriate controls. The urinary excretion of this biomarker was significantly associated with other parameters, including smoking habits, co-exposure to other airborne carcinogens and length of employment. In particular, multivariate regression analysis showed that the level of 15-F_{2t}-IsoP was linearly correlated to the length of exposure. In

agreement with our findings, other studies have previously reported increased oxidative stress generation in relation to occupational exposure to wood dust (Bruschweiler et al., 2016; Palus et al., 1999; Rekhadevi et al., 2009). Our findings provide strengthening of the hypothesis that oxidative stress and LPO can have a main role in the toxicity of wood dust. The analysis of F₂-IsoP in urine could offer a unique noninvasive analytic tool to study the role of ROS in chronic occupational exposures. In the current case, the linkage between urinary 15-F_{2t}-IsoPs and wood dust can be due to an increased production of ROS caused by inflammation after exposure fine and abundant airborne dust created during wood manipulation, maintenance activities and cleaning equipment. Increased oxidative stress and LPO can be caused from the oxidative burst of activated macrophages and neutrophils, cells with a main role in phagocytosis and clearance of xenobiotic particles, and from increased inflammatory cytokines and activated leukocytes (Gungor et al., 2010; Vanhees et al., 2013). This is in keeping with the results of previous studies using a biomarker of oxidative DNA damage and LPO (Bonassi et al., 2017; Bono et al., 2016; Bono et al., 2010; Peluso et al., 2013; Peluso et al., 2010). In support of our hypothesis, free radicals produced through chronic inflammatory process and cancer disease have been implicated as the causal factor in the mutagenesis of the *tumor suppressor gene TP53* (Brancato et al., 2016; Perez-Escuredo et al., 2012).

Next, our study showed an empirical relationship between tobacco smoking and the urinary excretion of 15-F_{2t}-IsoP, possibly related to the inhalation exposure to carcinogens contained in tobacco smoke. A 43% increment of the level of 15-F_{2t}-IsoP was present in overall the smokers as compared to the non-smokers. This excess is commonly interpreted as an harmful oxidative stress (Basu, 2008). These findings were somewhat expected as active smokers inhale a broad range of airborne carcinogens (IARC, 2004). The involvement of altered oxidative stress-related mechanisms in tobacco smoke carcinogenesis is in line with previous studies using various biomarkers of oxidative stress and LPO (Munnia et al., 2004; Peluso et al., 2014; Romanazzi et al., 2013). Various groups have measured the concentrations of F₂-IsoP in biological fluids of smokers. The mean level of free and esterified F₂-IsoP in the urine and plasma of smokers have been found to be significantly elevated as compared to non-smokers (Lowe et al., 2013). For instance, a previous cross-sectional study conducted on workers employed in an industry of plastic laminates in Piedmont, Italy, founds that smoking habits were significantly associated with the urinary excretion of 15-F_{2t}-IsoP (Romanazzi et al., 2013). When the relationship of 15-F_{2t}-IsoP with BMI was investigated, we found that the levels of 15-F_{2t}-IsoP tended to increase with fat accumulations. The 42% of the obese subjects showed indeed higher excretion of 15-F_{2t}-IsoP in respect to those with normal weight. This is partially in keeping with a previous work of Annor et al. (Annor et al., 2017) on the risk of diabetes and weight gain. In that study, the 35% of the obese individuals showed greater levels of F₂-IsoPs as compared to the controls. Additional studies are necessary to understand if this biomarker can be used as measure of lifestyle habits and intervention targeted to obesity prevention.

The threshold exposure limit recommended by the Italian law is 5 mg/m³ (Legislative Decree No 66/2000). This value will remain until the 2020th, after the entry into force of the new threshold exposure limit of 3 mg/m³ for five years and thereafter of 2 mg/m³ (European Directive Decree No 2017/2398). In this context, the SIREP database aims to facilitate analysis of occupational exposure figures for carcinogenic agents. In our study, the average amount of wood dust concentrations experienced from the wood workers was lower than threshold exposure limit of 3 mg/m³ (i.e., 1.48 mg/m³). This result is consistent with that reported from a previous study of Scarselli et al. (Scarselli et al., 2008), where the mean concentrations of wood dust was of 1.44 mg/m³ for 1.181 companies in Italy. Although our static measurements of the concentrations of industrial contaminants by fixed positions provide evidence of wood workers' exposure via air, they are not well representative of individual exposures to wood dust due to spatial and temporal variations. Therefore, we could not assess the potential relationships of airborne measurements with biomarker urinary excretion in exposed workers.

The airborne wood-dust concentrations from exposure registries are commonly used for the purposes of hazard control, exposure surveillance and assessment of health risks (Kauppinen et

al., 2006). Nevertheless, a limitation of our study is that no data on the variability of wood dust concentrations within a facility were available. The bias due to the variability of airborne carcinogen levels in occupational settings is difficult to predict, but a large variation can be present in one spot of a factory versus another. There could be an underestimation of the exposure to wood dust associated to some woodworking operations. For instance, local exhaust ventilation is used widely with fixed woodworking machinery, but it is generally lacking for hand tools (Pisaniello et al., 1991). The effects of poor work practices, such as the use of compressed air for cleaning, the lack of local exhaust ventilation for hand tools, that are commonly associated to high exposure levels to wood dust (Alwis et al., 1999), could be missed. Variations in the use of PPE (Alwis et al., 1999) and in the effective application of WorkSafe procedures at work places could have influenced the personal levels of exposure to wood dust of our workers.

Our subsequent finding shows that the urinary excretion of 15-F_{2t}-IsoP in the workers exposed to wood dust can aggravate with co-exposure to other respiratory carcinogens. An excess of 41% was detected in the wood workers that were co-exposed to organic solvents compared to the controls. Conversely, a lower excess was determined in the only wood dust exposed workers. High biosynthesis of F₂-IsoP can be due to frequent free radical-catalyzed reactions induced by alterations of oxidative stress, antioxidant defence and inflammation especially caused by occupational exposures to complex mixtures of airborne carcinogens. This is consistent with a cross-sectional study of workers exposed to dust containing silica (Peluso et al., 2015). In this study, the amount of oxidative stress and LPO biomarker of the workers exposed to airborne silica dust was greater in the case of occupational co-exposures to organic solvents, welding and motor exhaust fumes. Constituents of organic solvents, such as benzene and formaldehyde can be involved in the generation of oxidative stress and ROS (Bono et al., 2016; Bono et al., 2010; Sorensen et al., 2003) and cause the production of 15-F_{2t}-IsoP determined in the workers exposed to wood dust. Our results suggest that the urinary level of F₂-IsoP resulting from exposures to airborne wood dust can be affected from concomitant carcinogen exposures. Levels of oxidative stress can increase with exposures to organic solvents (Salimi et al., 2017; Singh et al., 2010), leading to a greater imbalance between excessive ROS generation and their degradation by antioxidants. The induction of reactive species can increase damage to membrane lipids, cellular proteins and DNA.

A significant difference in the amount of 15-F_{2t}-IsoP was then observed among sub-groups of wood workers with different occupational history. The urinary excretion of this biomarker of oxidant status was significantly elevated in those subjects with longer exposure time. An 41% excess of 15-F_{2t}-IsoP was found in the long-term wood workers as compared to those with shorter exposures, used as the reference level. Multivariate regression analysis showed that the level of 15-F_{2t}-IsoP was significantly linearly correlated to the length of employment, in agreement with a previous study on asbestos workers (Yoshida et al., 2001). In that study, the generation of an urinary biomarker of oxidative stress correlated positively with the length of exposure. Rekhadevi et al. (Rekhadevi et al., 2009) have similarly found an association between length of occupational exposure and increase frequency of micronuclei. Taken together, the occurrence of elevated oxidative stress in long-term wood workers can be possibly due to chronic inflammatory conditions. Our study suggests that the measure of urinary F₂-IsoPs can serve as a biomarker for assessing occupational carcinogen exposure and improving workplace safety. Particular effort should be devoted to studying long term health effects of exposure to wood dust, such as SNC.

Particular effort should be devoted to study delayed reactions such as diseases that take a long time to develop, like SNC, that can be caused by long-term exposure to this carcinogenic agent.

5. Conclusions

Our study provides a valuable contribution to the issue of oxidative stress in woodworking. An excessive ROS generation was demonstrated in exposed workers. Furthermore, we showed that exposure to organic solvents can increase the levels of urinary biomarkers of oxidative stress in wood workers. Results provide a basis for worker surveillance in occupational settings. F₂-IsoP measure could be used for the evaluation of the effectiveness of targeted interventions aimed to

356 reduce exposures to various environmental carcinogens. A more effective control of occupational
357 health risks could decrease the incidence of illness at work and improve the health of the
358 workforce.

359 **Acknowledgments**

360 We are grateful to Dr. Dusca Bartoli, Dr. Giuseppe A. Farina, Dr. Tonina E. Iaia, Dr. Pierluigi Faina,
361 Dr. Luciano Monticelli for the assistances contacting with workers.

362 **Funding**

363 This work was supported by the Tuscany Region and the National Institute for Insurance Against
364 Accident at Work (INAIL).

365 **Declarations of interest**

366 None

367

References

- Acheson, E. D., et al., 1968. Nasal cancer in woodworkers in the furniture industry. *Br Med J.* 2, 587-596.
- Alwis, U., et al., 1999. Dust Exposures in the Wood Processing Industry. *Am Ind Hyg Ass J.* 60, 641-646.
- Annor, F., et al., 2017. African Ancestry Gradient Is Associated with Lower Systemic F(2)-Isoprostane Levels. *Oxid Med Cell Longev.* 2017, 8319176-8319176.
- Ball, M. J., 1968. Nasal cancer in woodworkers. *Br Med J.* 3, 253-253.
- Bartels, H., Cikes, M., 1969. [Chromogens in the creatinine determination of Jaffe]. *Clin Chim Acta.* 26, 1-10.
- Basu, S., 2008. F2-isoprostanes in human health and diseases: from molecular mechanisms to clinical implications. *Antiox Redox Sign.* 10, 1405-1434.
- Binazzi, A., et al., 2017. Sinonasal cancer in the Italian national surveillance system: Epidemiology, occupation, and public health implications. *Am J Ind Med.* 1-12.
- Bonassi, S., et al., 2017. 3-(2-deoxy- β -d-erythro-penta-furanosyl) pyrimido [1, 2- α] purin-10 (3H)-one deoxyguanosine adducts of workers exposed to asbestos fibers. *Toxic Lett.* 270, 1-7.
- Bono, R., et al., 2016. Formaldehyde-induced toxicity in the nasal epithelia of workers of a plastic laminate plant. *Toxicol Res.* 5, 752-760.
- Bono, R., et al., 2010. Malondialdehyde-deoxyguanosine adduct formation in workers of pathology wards: the role of air formaldehyde exposure. *Chem Res Toxicol.* 23, 1342-8.
- Bono, R., et al., 2015. Urban air and tobacco smoke as conditions that increase the risk of oxidative stress and respiratory response in youth. *Environ Res.* 137, 141-146.
- Brancato, B., et al., 2016. 8-Oxo-7, 8-dihydro-2-deoxyguanosine and other lesions along the coding strand of the exon 5 of the tumour suppressor gene P53 in a breast cancer case-control study. *DNA Res.* 23, 395-402.
- Bruschweiler, D. E., et al., 2016. DNA Damage among Wood Workers Assessed with the Comet Assay. *Environ Health Insights.* 10, 105-112.
- Bruschweiler, E. D., et al., 2014. Workers exposed to wood dust have an increased micronucleus frequency in nasal and buccal cells: results from a pilot study. *Mutagenesis.* 29, 201-207.
- Çelik, A., Kanık, A., 2006. Genotoxicity of occupational exposure to wood dust: Micronucleus frequency and nuclear changes in exfoliated buccal mucosa cells. *Environ Mol Mutagen.* 47, 693-698.
- Ceppi, M., et al., 2011. Study design and statistical analysis of data in human population studies with the micronucleus assay. *Mutagenesis.* 26, 247-252.
- d'Errico, A., et al., 2009. A case-control study on occupational risk factors for sino-nasal cancer. *Occup Environ Med.* 66, 448-455.
- Demers, P. A., et al., 1995. Wood dust and sino-nasal cancer: pooled reanalysis of twelve case-control studies. *Am J Ind Med.* 28, 151-66.
- Gungor, N., et al., 2010. Transcriptional profiling of the acute pulmonary inflammatory response induced by LPS: role of neutrophils. *Respir Res.* 11, 24.
- Hancock, D. G., et al., 2015. Wood dust exposure and lung cancer risk: a meta-analysis. *Occup Environ Med.* 72, 889-898.
- IARC, 1995. Wood Dust and Formaldehyde. *IARC Monogr Eval Carcinog Risk Chem Hum.* 62, 35-215.
- IARC, 2004. Tobacco smoke and involuntary smoking. *IARC Monogr Eval Carcinog Risks Hum.* 83, 1-1438.
- IARC, 2012. Wood Dust. A Review of Human Carcinogens: Arsenic, Metals, Fibres, and Dusts. *IARC Monogr Eval Carcinog Risks Hum.* 100 C, 1-469.
- Il'yasova, D., et al., 2015. Urinary F2-Isoprostanes and Metabolic Markers of Fat Oxidation. *Oxid Med Cell Long.* 2015.
- Janicka, M., et al., 2010. Isoprostanes-Biomarkers of Lipid Peroxidation: Their Utility in Evaluating Oxidative Stress and Analysis. *IJMS.* 11, 4631.
- Kauppinen, T., et al., 2006. Occupational Exposure to Inhalable Wood Dust in the Member States of the European Union. *Ann Occ Hyg.* 50, 549-561.
- Lowe, F. J., et al., 2013. Lung cancer biomarkers for the assessment of modified risk tobacco products: an oxidative stress perspective. *Biomarkers.* 18, 183-195.
- Marnett, L. J., 2000. Oxyradicals and DNA damage. *Carcinogenesis.* 21, 361-70.

- Merlo, F., et al., 1997. Airborne levels of polycyclic aromatic hydrocarbons: 32P-postlabeling DNA adducts and micronuclei in white blood cells from traffic police workers and urban residents. *J Environ Pathol Toxicol Oncol.* 16, 157-62.
- Milne, L. G., et al., 2015. The isoprostanes—25 years later. *BBA-Mol Cell Biol L.* 1851, 433-445.
- Montuschi, P., et al., 2004. Isoprostanes: markers and mediators of oxidative stress. *FASEB.* 18, 1791-1800.
- Morrow, J. D., et al., 1999. Quantification of the Major Urinary Metabolite of 15-F₂t-Isoprostane (8-iso-PGF₂±) by a Stable Isotope Dilution Mass Spectrometric Assay. *Anal Bioch.* 269, 326-331.
- Munnia, A., et al., 2004. Exocyclic malondialdehyde and aromatic DNA adducts in larynx tissues. *Free Radic Biol Med.* 37, 850-8.
- Munnia, A., et al., 2017. Bulky DNA Adducts, Tobacco Smoking, Genetic Susceptibility, and Lung Cancer Risk. *Adv Clin Chem.* 81, 231-77.
- Munnia, A., et al., 2007. 32P-Post-labelling method improvements for aromatic compound-related molecular epidemiology studies. *Mutagenesis.* 22, 381-5.
- Palus, J., et al., 1999. DNA damage detected by the comet assay in the white blood cells of workers in a wooden furniture plant. *Mutat Res.* 444, 61-74.
- Peluso, M., et al., 1997. Detection of DNA adducts in human nasal mucosa tissue by 32P-postlabeling analysis. *Carcinogenesis.* 18, 339-44.
- Peluso, M., et al., 2012. DNA methylation differences in exposed workers and nearby residents of the Ma Ta Phut industrial estate, Rayong, Thailand. *Int J Epidemiol.* 41, 1753-60; author response 1761-3.
- Peluso, M., et al., 2013. Malondialdehyde-deoxyguanosine and bulky DNA adducts in schoolchildren resident in the proximity of the Sarroch industrial estate on Sardinia Island, Italy. *Mutagenesis.* 28, 315-21.
- Peluso, M., et al., 2010. Malondialdehyde-deoxyguanosine adducts among workers of a Thai industrial estate and nearby residents. *Environ Health Perspect.* 118, 55-9.
- Peluso, M. E., et al., 2014. Aberrant methylation of hypermethylated-in-cancer-1 and exocyclic DNA adducts in tobacco smokers. *Toxicol Sci.* 137, 47-54.
- Peluso, M. E., et al., 2015. Oxidatively damaged DNA in the nasal epithelium of workers occupationally exposed to silica dust in Tuscany region, Italy. *Mutagenesis.* 30, 519-25.
- Perez-Escuredo, J., et al., 2012. Wood dust-related mutational profile of TP53 in intestinal-type sinonasal adenocarcinoma. *Human Pathology.* 43, 1894-1901.
- Pisaniello, D. L., et al., 1991. Wood dust exposure during furniture manufacture -results from an Australian survey and consideration for threshold limit value development. *Am Ind Hyg Ass J.* 52, 485-492.
- Pylkkänen, L., et al., 2009. Wood dusts induce the production of reactive oxygen species and caspase-3 activity in human bronchial epithelial cells. *Toxic in Vitro.* 262, 265-270.
- Rekhadevi, P. V., et al., 2009. Genetic damage in wood dust-exposed workers. *Mutagenesis.* 24, 59-65.
- Roberts, L. J., Morrow, J. D., 2000. Measurement of F₂-isoprostanes as an index of oxidative stress in vivo. *Free Radic Biol Med.* 28, 505-513.
- Romanazzi, V., et al., 2013. 15-F₂t isoprostane as biomarker of oxidative stress induced by tobacco smoke and occupational exposure to formaldehyde in workers of plastic laminates. *Sci Total Environ.* 442, 20-5.
- Salimi, A., et al., 2017. Xylene Induces Oxidative Stress and Mitochondria Damage in Isolated Human Lymphocytes. *Toxic res.* 33, 233-238.
- Scarselli, A., et al., 2008. Occupational exposure levels to wood dust in Italy, 1996–2006. *Occup Environ Med.* 65, 567-574.
- Singh, M. P., et al., 2010. Effects of co-exposure of benzene, toluene and xylene to *Drosophila melanogaster*: Alteration in hsp70, hsp60, hsp83, hsp26, ROS generation and oxidative stress markers. *Chemosphere.* 79, 577-587.
- Sorensen, M., et al., 2003. Linking exposure to environmental pollutants with biological effects. *Mutat Res.* 544, 255-71.
- van Houwelingen, H. C., et al., 2002. Advanced methods in meta-analysis: multivariate approach and meta-regression. *Stat Med.* 21, 589-624.

- Vanhees, K., et al., 2013. Intrauterine exposure to flavonoids modifies antioxidant status at adulthood and decreases oxidative stress-induced DNA damage. *Free Radic Biol Med.* 57, 154-61.
- Wultsch, G., et al., 2015. Impact of exposure to wood dust on genotoxicity and cytotoxicity in exfoliated buccal and nasal cells. *Mutagenesis.* 30, 701-709.
- Yoshida, R., et al., 2001. Urinary 8-oxo-7, 8-dihydro-2'-deoxyguanosine and biopyrrins levels among construction workers with asbestos exposure history. *Ind Health.* 39, 186-8.

Table 1. Demographics and other variables by exposure group.

	Study Population		
	Controls	Wood workers	
	N	N	N
Study population	57	123	
Age (years)	47.2 ± 11	44.4 ± 11	
Smoking habits			
Non-smokers	26	56	
Ex-smokers	12	23	
Smokers	19	44	
Body mass index (BMI)	25 ± 0.40	25 ± 0.27	
BMI categories			
Normal weight (18.5-24.9)	27	67	
Overweight (25-24.99)	28	46	
Obese (≥30)	2	10	

Table 2. Mean level of 15-F_{2t} isoprostane (15-F_{2t}-IsoP) and Mean Ratio (MR) and 95% Confidence Interval (C.I.) by exposure group and other variables.

Urinary 15-F_{2t} Isoprostane and Wood Dust Exposure				
	N	15-F_{2t} IsoP ± SE	MR, 95% C.I.	P-value^a
Smoking habits				
Non-smokers	82	3.3 ± 0.19	Reference level	
Ex-smokers	35	3.7 ± 0.38	1.15, 0.96-1.01	0.092
Smokers	63	4.5 ± 0.31	1.43, 1.23-1.66	<0.0001
Exposure group				
Controls	57	2.9 ± 0.19	Reference level	
Wood workers	123	4.2 ± 0.21	1.36, 1.18-1.57	<0.0001
Co-carcinogen occupational exposures				
Controls	57	2.9 ± 0.19	Reference level	
Only wood dust exposed workers	85	4.1 ± 0.25	1.34, 1.15-1.56	0.0001
Wood dust with organic solvents and formaldehyde exposed workers	38	4.5 ± 0.43	1.41, 1.17-1.70	0.0002
Occupational history				
≤8 years	38	3.2 ± 0.18	Reference level	
9-25 years	43	4.4 ± 0.38	1.27, 1.04-1.55	0.017
≥26 years	42	4.8 ± 0.44	1.41, 1.14-1.75	0.0014
Body mass index categories				
Normal weight (18.5-24.9 kg/m ²)	94	3.6 ± 0.16	Reference level	
Overweight (25-24.99 kg/m ²)	74	4.0 ± 0.32	1.05, 0.90-1.21	0.5393
Obese (≥30 kg/m ²)	12	4.4 ± 0.84	1.10, 0.84-1.44	0.5018

^a P-values (Test of Wald) were adjusted for age and smoking, as appropriate.

Table 3. Average level of 15-F_{2t} isoprostane (15-F_{2t}-IsoP) and Mean Ratio (MR) and 95% Confidence Interval (C.I.) by exposure group after smoking stratification.

Excess Risk of Urinary Biomarker in Wood Dust Workers				
	15-F_{2t}-IsoP ± SE	MR, 95% C.I.	P-value ^a	
	N	ng/mg creatinine		
Non-smokers				
Controls	26	2.8 ± 0.37	Reference level	
Wood workers	56	3.5 ± 0.23	1.27, 1.01-1.59	0.0353
Ex-smokers				
Controls	12	2.7 ± 0.30	Reference level	
Wood workers	23	4.3 ± 0.53	1.48, 1.09-2.01	0.01
Smokers				
Controls	19	3.3 ± 0.17	Reference level	
Wood workers	44	5.0 ± 0.42	1.53, 1.23-1.91	0.0001

^a P-values (Test of Wald) were adjusted for age.